

Weekly Influenza

SURVEILLANCE REPORT

Disease Week 50 Highlights 12/11/2022-12/17/2022


Influenza

- Influenza cases continue to remain high but are declining with only 183 cases reported last week.
- Four influenza-associated deaths were reported this week, for a total of 6 influenza-associated deaths reported so far this season.
- The average age of influenza-associated deaths was 69 years and ranged between 62 years and 83 years.
- A total of six influenza outbreaks have been reported to date, with one new outbreak in a long-term care facility (LTCF).
- Influenza-like illness activity decreased slightly during Week 49. The reported percentage of visits for ILI decreased to 16.6% compared to 25.5% reported the previous week.

Respiratory syncytial virus (RSV)

- RSV cases appeared to have peaked in mid-November and are now declining.
- Five new RSV cases were reported this week, all among children less than 5 years, bringing the total to 116 cases.

City of Long Beach Department of Health and Human Services Epidemiology Program




*This report was revised in February 2023.




Prepared by the Department of Health and Human Services

OVERVIEW

Total Cases¹
2166
Outbreaks²
6
Deaths³





*Data for December 2022 is not complete for the month.

195 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183 | 183


- 1. Total case counts are based on those reported to public health, the true number of influenza cases are under reported.
- 2. Outbreaks are defined as at least one case of laboratory confirmed influenza and at least two residents with onset of influenza-like-illness (ILI) within 72 hours. If an outbreak is in the community setting (i.e. school or daycare), outbreak is defined as 5 or more cases of ILI within a group within 72 hours.
- 3. Number of deaths is based on influenza-coded deaths from death certificates. They are not necessarily laboratory-confirmed and may be an underestimate of all influenza-associated deaths.









Prepared by the Department of Health and Human Services






INFLUENZA BY RACE/ETHNICITY, 2022-2023



INFLU	ENZA AND PN	EUMONIA ⁵	
Season	Influenza Deaths	Pneumonia Deaths	% Influenza & Pneumonia Deaths
2019 - 2020	11	369	11.0%
2020 - 2021	0	589	14.7%
2021 - 2022	1	344	9.8%
2022 - 2023	6	40	8 1%



5. The number of influenza and pneumonia related deaths is based on causes of death listed on the death certificates. Deaths are not necessarily lab confirmed influenza or pneumonia.







Prepared by the Department of Health and Human Services

Influenza-Like Illness ED Encounters


Influenza-like Illness (ILI) emergency department (ED) encounters are based on syndromic surveillance data from one syndromic-participating hospital in Long Beach. Syndromic surveillance is a population-based symptom monitoring system that uses hospital-based data. This report presents ILI ED encounters from 2020 through the current influenza season (2022-2023). ILI was defined as emergency department encounters with a chief complaint mentioning influenza or fever and cough or fever and sore throat.

Please note that syndromic keywords and codes specific to ILI are broad enough to include other respiratory conditions including COVID-19.

Syndromic surveillance data will have a 1- week lag due to the current data transfer from the county.

INFLUENZA-LIKE	ILLNESS ED ENCOU	INTERS, WEEK 49
2020-2021	2021-2022	2022-2023
3.9%	4.0%	16.6%

INFLUENZA-LIKE ILLNESS ED ENCOUNTERS, 2020 – 2023 26,3% 21,1% 20% 10% 2,3% 1,1% 3,9% 3,6% 3,6% 5,4% 2,1% Jul 2020 Jan 2021 Jul 2021 Jul 2022 ED Encounter Week







Prepared by the Department of Health and Human Services

Respiratory Syncytial Virus Infection (RSV)

Respiratory syncytial virus, or RSV, is a common respiratory virus that usually causes mild, cold-like symptoms. Most people recover in a week or two, but RSV can be serious, especially for infants and older adults. RSV is the most common cause of bronchiolitis (inflammation of the small airways in the lung) and pneumonia (infection of the lungs) in children younger than 1 year of age in the United States. Clinical symptoms of RSV are nonspecific and can overlap with other viral respiratory infections, as well as some bacterial infections.

TOTAL 2022-2023 CASES


116

NEW WEEKLY CASES


5

PEDIATRIC DEATHS

0



*Data for December 2022 is not complete for the month.









Prepared by the Department of Health and Human Services

RSV BY AGE AND SEASON							
Age	2018-2019	2019-2020	2020-2021	2021-2022	2022-2023		
0-4	68%	86%	0%	68%	81%		
05-17	0%	3%	0%	4%	10%		
18-64	14%	3%	0%	14%	5%		
65+	18%	8%	0%	14%	3%		



