



MPOX VACCINE

What to Expect After Getting Vaccinated

Common Side Effects to Expect

If the vaccine was given intradermally, some people have reported:

- Itching
- Swelling
- Redness
- Thickening of the skin
- Skin discoloration at the site of injection



Some of these side effects may last for several weeks. If you have concerns about receiving the vaccine intradermally, you can ask for the vaccine to be given to you subcutaneously in the fat layer underneath the skin on the back of your upper arm (triceps).

It is important to take extra caution if you have had an allergic reaction to any vaccine. You may still be vaccinated with JYNNEOS, but your provider may need to observe you for 30 minutes after you get vaccinated to make sure you don't develop an allergic reaction. Talk to your healthcare provider if you had an allergic reaction to the **antibiotics gentamicin or ciprofloxacin, or chicken or egg protein.**

Severe allergic reaction may include:



Hives



Swollen Face



Difficulty
Breathing



Fast Heartbeat



Dizziness



Weakness

If you think you are having a severe allergic reaction, call 911 or go to the hospital. Call your health care provider if you have other side effects that concern you or call 311 to be connected to care.

Your provider is required to report serious adverse reactions to the Vaccine Adverse Event Reporting System (VAERS), however you can report as well. VAERS staff members do not give medical advice

To Report VAERS



vaers.hhs.gov



800-822-7967



MPOX VACCINE


What to Expect After Getting Vaccinated

Mpox Prevention

- Avoid close, skin-to-skin contact with the mpox rash.
- Do not touch the rash or scabs of a person with mpox.
- Do not kiss, hug, cuddle, or have sex with someone with mpox.
- Do not share eating utensils or cups.
- Do not handle or touch the bedding, towels, or clothing of a sick person.
- Wash your hands often with soap and water or use an alcohol-based hand sanitizer, especially after contact with sick people.
- Some people may experience side effects more severely and/or for a longer duration. Please talk to a health care professional if you do not feel well after 2 or 3 days.

VACCINE REMINDERS

- If the Jynneos vaccine is given before exposure or within 4 days of exposure, Jynneos may reduce the chance of infection. If given between 5 and 14 days after exposure, it may lower the severity of symptoms.
- You will not be fully vaccinated and getting the most benefit from this vaccine until 2 weeks after your second shot of JYNNEOS.
- The second dose should be given 4 weeks after the first dose for maximum protection.



For more information:

For additional questions regarding side effects contact your health care provider, or refer JYNNEOS package insert



562.570.7912



longbeach.gov/mpox